

Lung volume reduction surgery for advanced emphysema

Interventional procedures guidance

Published: 23 February 2005

[nice.org.uk/guidance/ipg114](https://www.nice.org.uk/guidance/ipg114)

Your responsibility

This guidance represents the view of NICE, arrived at after careful consideration of the evidence available. When exercising their judgement, healthcare professionals are expected to take this guidance fully into account. However, the guidance does not override the individual responsibility of healthcare professionals to make decisions appropriate to the circumstances of the individual patient, in consultation with the patient and/or guardian or carer.

Commissioners and/or providers have a responsibility to implement the guidance, in their local context, in light of their duties to have due regard to the need to eliminate unlawful discrimination, advance equality of opportunity, and foster good relations. Nothing in this guidance should be interpreted in a way that would be inconsistent with compliance with those duties.

Commissioners and providers have a responsibility to promote an environmentally sustainable health and care system and should assess and reduce the environmental impact of implementing NICE recommendations wherever possible.

1 Guidance

- 1.1 Current evidence on the safety and efficacy of lung volume reduction surgery for advanced emphysema appears adequate to support the use of this procedure provided that the normal arrangements are in place for consent, audit and clinical governance.

- 1.2 Clinicians wishing to use lung volume reduction surgery for advanced emphysema should ensure that patients are fully informed about the risks of the procedure and the likelihood of deterioration in the longer term. Use of the Institute's [information for the public](#) is recommended.
- 1.3 Patient selection is important because mortality is increased in patients with the most seriously compromised lung function. The Institute has issued a clinical guideline on [chronic obstructive pulmonary disease](#).
- 1.4 The procedure should be undertaken by a multidisciplinary team that includes a respiratory physician, specialists in pulmonary rehabilitation and a thoracic surgeon.

2 The procedure

2.1 *Indications*

- 2.1.1 Emphysema is a chronic lung disease. The walls of the air sacs (alveoli) in the lung weaken and disintegrate, leaving behind abnormally large air spaces that remain filled with air even when the patient breathes out. These air spaces may coalesce to form larger air-filled sacs called bullae. The surface area of the alveoli is decreased, so there is less space for the exchange of oxygen and carbon dioxide. This leads to reduced levels of oxygen in the blood. The most common symptoms of emphysema are shortness of breath (dyspnoea), coughing, fatigue and weight loss.
- 2.1.2 Emphysema often co-exists with chronic bronchitis. Both of these conditions may be described by the more general term of chronic obstructive pulmonary disease (COPD).
- 2.1.3 Treatment for COPD involves a multidisciplinary approach, which may include education, exercise, breathing retraining, smoking cessation, oral and inhaled medication, oxygen therapy, and lung transplantation. Lung volume reduction surgery may be an option for patients with severe symptoms for whom conservative treatments have proved inadequate.

2.2 *Outline of the procedure*

- 2.2.1 Lung volume reduction surgery is a palliative treatment that aims to remove the least functional part of the lungs. Computed tomography (CT) and perfusion scanning are used to identify the diseased lung tissue. The diseased part of the lung can be accessed by various techniques including median sternotomy, video-assisted thoracoscopy and thoracotomy. The first two are the most common techniques. Median sternotomy involves cutting through the sternum to open the chest. The video-assisted procedure involves making a number of small incisions in both sides of the chest to allow the insertion of instruments into the chest between the ribs. A thoracotomy involves making an incision between the ribs on one side of the chest and separating the ribs to access the lung.
- 2.2.2 The aim of the surgery is to reduce the volume of the lung. This is done by using a surgical stapling device to cut and seal the tissue, laser ablation to shrink lung volume, or a combination of both. Once the tissue has been removed, the lung is re-inflated and the chest closed.

2.3 *Efficacy*

- 2.3.1 Evidence on efficacy indicates that in certain patients lung function, exercise performance and quality of life are improved in the short term after lung volume reduction surgery. These results have been relatively consistent across study designs and were confirmed in the National Emphysema Treatment Trial, a recent large-scale randomised controlled trial comparing surgery with medical therapy.
- 2.3.2 The National Emphysema Treatment Trial randomised 1218 patients, of whom 580 underwent surgery. At 24 months, exercise capacity had improved in 15% (54/371) of patients in the surgery group compared with 3% (10/378) of patients in the medical group ($p < 0.001$). Quality of life had also improved in the surgical group (121/371) as compared with the medical group (34/378) at 24 months (33% versus 9%, $p < 0.001$). However, the trial found no difference in overall mortality between the two groups (0.11 deaths per person-year, risk ratio 1.01, $p = 0.90$). For more details, refer to the Sources of evidence section.
- 2.3.3 The Specialist Advisors considered that the procedure is beneficial for a select proportion of patients, but the benefit tends to decline with time.

2.4 *Safety*

- 2.4.1 The most common complication was persistent air leak from the lung. In one study of 250 patients, 45% (113/250) of patients experienced prolonged air leaks lasting more than 7 days, with 8 of these patients (3%) requiring a subsequent operation. Other complications in this series included pneumonia 10% (24/250), in-hospital mortality 5% (12/250), myocardial infarction 2% (5/250), deep vein thrombosis 2% (4/250), small bowel obstruction 2% (6/250) and phrenic nerve injury < 1% (2/250). For more details, refer to the Sources of evidence section.
- 2.4.2 Complications include those that may arise from pre-existing co-morbidities as well as those that are directly due to the surgery.
- 2.4.3 The Specialist Advisors considered that the risks of surgery were well known. They listed the main complications as being air leaks, chest infections and respiratory failure.

2.5 *Other comments*

- 2.5.1 It was noted that endobronchial techniques are being used increasingly as an alternative to this procedure.

Andrew Dillon
Chief Executive
February 2005

3 Further information

Sources of evidence

The evidence considered by the Interventional Procedures Advisory Committee is described in the following document.

'Interventional procedures overview of lung volume reduction surgery for advanced emphysema', June 2004.

Information for patients

NICE has produced [information on this procedure for patients and carers](#) ('Understanding NICE guidance'). It explains the nature of the procedure and the guidance issued by NICE, and has been written with patient consent in mind.

4 About this guidance

NICE interventional procedure guidance makes recommendations on the safety and efficacy of the procedure. It does not cover whether or not the NHS should fund a procedure. Funding decisions are taken by local NHS bodies after considering the clinical effectiveness of the procedure and whether it represents value for money for the NHS. It is for healthcare professionals and people using the NHS in England, Wales, Scotland and Northern Ireland, and is endorsed by Healthcare Improvement Scotland for implementation by NHSScotland.

This guidance was developed using the NICE [interventional procedure guidance](#) process.

This guidance has been incorporated into the [NICE pathway on chronic obstructive pulmonary disease](#), along with other related guidance and products.

We have produced a [summary of this guidance for patients and carers](#). Information about the evidence it is based on is also [available](#).

Changes since publication

24 January 2012: minor maintenance.

Your responsibility

This guidance represents the views of NICE and was arrived at after careful consideration of the available evidence. Healthcare professionals are expected to take it fully into account when exercising their clinical judgement. This guidance does not, however, override the individual responsibility of healthcare professionals to make appropriate decisions in the circumstances of the individual patient, in consultation with the patient and/or guardian or carer.

Implementation of this guidance is the responsibility of local commissioners and/or providers. Commissioners and providers are reminded that it is their responsibility to implement the guidance, in their local context, in light of their duties to avoid unlawful discrimination and to have

regard to promoting equality of opportunity. Nothing in this guidance should be interpreted in a way which would be inconsistent with compliance with those duties.

Copyright

© National Institute for Health and Clinical Excellence 2005. All rights reserved. NICE copyright material can be downloaded for private research and study, and may be reproduced for educational and not-for-profit purposes. No reproduction by or for commercial organisations, or for commercial purposes, is allowed without the written permission of NICE.

Contact NICE

National Institute for Health and Clinical Excellence
Level 1A, City Tower, Piccadilly Plaza, Manchester M1 4BT

www.nice.org.uk

nice@nice.org.uk

0845 033 7780

Endorsing organisation

This guidance has been endorsed by [Healthcare Improvement Scotland](#).